

Math 106 Winter 2015

Test 2 (50 points)

Name: _____

Show all your work to receive full credit for a problem. Points will be taken off if you do not show how you arrived at your answer, even if the final answer is correct.

Please keep your written answers brief; be clear and to the point. Points will be taken off for rambling and for incorrect or irrelevant statements.

Do not use the calculator integral function. Whenever possible, find the exact values of integrals by finding antiderivatives or using the table of integrals.

When you use a formula from the table of integrals, mention the formula number and the value(s) of any constant(s) that you may need.

Give exact answers. If needed, round off your answers to four decimal places.

Include units in your answers wherever possible.

There are seven questions. Questions are printed on both sides of a page.

You may use any of the following facts:

$$P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \cdots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$$

$$|f(x) - P_n(x)| \leq \frac{K_{n+1}}{(n+1)!} |x - x_0|^{n+1}$$

$$\int u dv = uv - \int v du$$

$$\int_1^{\infty} \frac{1}{x^p} dx \text{ converges for } p > 1 \text{ and diverges for } p \leq 1.$$

$$\int_0^{\infty} e^{-ax} dx \text{ converges for } a > 0.$$

1. (8 points) Evaluate the following integral. (You may use formulas 1-18, 39-42, 50, 51 only from the table of integrals for this problem.)

$$\int x^2 \cos x \, dx$$

2. (7 points) Evaluate the following integral. (You may use formulas 1-18, 39-42, 50, 51 only from the table of integrals for this problem.)

$$\int \tan^3(2x) \sec^4(2x) dx$$

3. (8 points) Evaluate the following integral. (You may use formulas 1-18, 39-42, 50, 51 only from the table of integrals for this problem.)

$$\int \sqrt{64 + x^2} \, dx$$

4. (8 points) Evaluate the following integral. (You may use formulas 1-18 only from the table of integrals for this problem.)

$$\int \frac{11 + 3x + 2x^2}{(x^2 + 16)(x - 7)} dx$$

5. (5 points) Use comparison to determine the convergence of the following integral.

$$\int_8^{\infty} \frac{2 \cos x + 3}{x^4 + 7} dx$$

6. (7 points) The p.d.f. of a continuous random variable X is given by $f(x) = ke^{-0.2x}$ for $x \geq 0$ (the function is zero for all other values of x). Find k .

7. (7 points) Suppose the fourth-order Taylor polynomial for a function f based at $x_0 = 1$ is

$$P_4(x) = (x - 1) + 2(x - 1)^2 + 3(x - 1)^4.$$

(a) Write the third-order Taylor polynomial for f based at $x_0 = 1$.

(b) What does Taylor's theorem imply about the maximum approximation error committed by P_4 over the interval $[-2, 2]$? Assume that $-4 \leq f^{(5)}(x) \leq 2$ for all x in $[-2, 2]$.