

Name: Solutions

Math 105: Fall 2013
Exam 1 (version π): October 4

Correct answers accompanied by incorrect or incomplete work will not receive full credit.


1. (6 points) A portion of the graph of $g(x) = x^3 - 3x$ is shown

(a) Draw the line tangent to $g(x)$ at $x = -0.5$.

(b) Compute the slope of your tangent line.

$$(-0.5, 1.4), (-0.75, 2)$$

$$\frac{2 - 1.4}{-0.75 - (-0.5)} = \frac{0.6}{-0.25} = -2.4$$



2. (8 points) Let $g(x) = x^3 - 3x$. Use the limit definition of derivative to find $g'(-0.5)$.

(Hint: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.)

$$g'(-0.5) = \lim_{h \rightarrow 0} \frac{g(-0.5+h) - g(-0.5)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(-0.5+h)^3 - 3(-0.5+h) - [(-0.5)^3 - 3(-0.5)]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{(-0.5)^3} + 3(-0.5)^2h + 3(-0.5)h^2 + h^3 + 1.5 - 3h - \cancel{(-0.5)^3} + 1.5}{h}$$

$$= \lim_{h \rightarrow 0} \frac{0.75h - 1.5h^2 + h^3 - 3h}{h} = \lim_{h \rightarrow 0} (-2.25 - 1.5h + h^2)$$

$$= -2.25$$

3. (6 points) Let $g(x) = x^3 - 3x$. Use the power rule to find $g'(-0.5)$.

$$g'(x) = 3x^2 - 3 \rightarrow g'(-0.5) = 3(-0.5)^2 - 3 = -2.25$$


4. (4 points) Let $P(t)$ be the price (in dollars) of gas t days after the start of the government shut down. What does the statement $P'(20) = -0.28$ mean in this context? Include units in your answer.

20 days after the shut down the price of gas is decreasing at a rate of 0.28 \$/day.

5. (4 points each) The graph of $f(x)$ is given. Evaluate the following (assume the tickmarks occur at 1, 2, etc).


(a) $f(3) = -2$

(b) $\lim_{x \rightarrow 3} f(x) = 0$




6. (10 points) Let $k(x) = |x|$.

- (a) Find $\lim_{x \rightarrow 0} k(x)$. Justify your answer.



$\lim_{x \rightarrow 0} k(x) = 0$, we can see this by looking at the graph at left.

- (b) Find $\lim_{x \rightarrow 0} k'(x)$. Justify your answer.



I drew the graph of $k'(x)$ at left,

$\lim_{x \rightarrow 0^+} k'(x) = 1$
 $\lim_{x \rightarrow 0^-} k'(x) = -1$ } since these are not equal

$\lim_{x \rightarrow 0} k'(x)$ DNE

7. (14 points) Let $h(x) = -7x^5 + \frac{12}{x^5} + \sqrt[3]{x} + \sin 12$. $= -7x^5 + 12x^{-5} + x^{1/3} + \sin 12$

(a) Find the derivative of h .

$$h'(x) = -35x^4 - 60x^{-6} + \frac{1}{3}x^{-2/3}$$

(b) Find the antiderivative of h .


$$H(x) = \frac{-7x^6}{6} + \frac{12x^{-4}}{-4} + \frac{x^{4/3}}{4/3} + (\sin 12)x + C$$

$$= -\frac{7}{6}x^6 - 3x^{-4} + \frac{3}{4}x^{4/3} + (\sin 12)x + C$$

8. (14 points) Assume that f is a continuous function defined on the closed interval $[-3, 3]$ such that $f(-3) = 3$ and $f(3) = -1$. Furthermore, assume that f' and f'' are continuous on $(-3, 3)$ and that the information in the table below is known about these functions. On the grid below sketch f

| x | $-3 \leq x < -1$ | -1 | $-1 < x < 0$ | 0 | $0 < x < 1$ | 1 | $1 < x \leq 3$ |
|----------|------------------|------|--------------|-----|-------------|-----|----------------|
| $f'(x)$ | - | 0 | - | - | - | 0 | + |
| $f''(x)$ | + | 0 | - | 0 | + | + | + |

f | dec, cc↑ | terrace | dec, cc↓ | inf | dec, cc↑ | min | inc, cc↑




just one possibility is shown at left.

9. (8 points) Assume $y = -\frac{2}{3}x + 4$ is tangent to $f(x)$ at $x = 5$.

(a) Find $f(5)$. Justify your answer.

find the y-coordinate when $x = 5$

$$y = -\frac{2}{3}(5) + 4 = -\frac{10}{3} + \frac{12}{3} = \boxed{\frac{2}{3}}$$




(b) Find $f'(5)$. Justify your answer.

slope of tangent line equals derivative.

so $\boxed{-\frac{2}{3}}$ which is the slope of the line

10. (18 points) The graph below is a graph of $g(x)$. Let $G(x)$ be an antiderivative of $g(x)$.



(a) Is it possible that $g''(1.5) = 2$? Justify your answer.

No, b/c g is concave down at $x = 1.5$ and the 2nd derivative of g is negative when g is concave down

(b) Is $G(0) > G(1)$? Justify your answer.

$G' = g$, since $g(x)$ is positive for all $0 \leq x \leq 1$, the slope of G is positive, hence G is increasing on the interval $[0, 1]$. So no, instead $G(0) < G(1)$.

(c) On what interval(s) is $G(x)$ concave up? Justify your answer.

$G(x)$ is cc \uparrow when $G' = g$ is increasing.

So $(-0.5, 1.5) \cup (4, \infty)$

11. (4 points) Who do you think will win the World Series?

- (a) Atlanta Braves (b) Boston Red Sox (c) Detroit Tigers (d) Los Angeles Dodgers
 (e) Oakland Athletics (f) Pittsburgh Pirates (g) St. Louis Cardinals (h) Tampa Bay Rays