

Name: Solutions

Math 105B: Fall 2012

Exam 1: October 5


Correct answers accompanied by incorrect or incomplete work will not receive full credit.

1. (8 points) Let $g(x) = b^x$ where b is a constant and $0 < b < 1$.

(a) What is the domain of $g(x)$?

$(-\infty, \infty)$

note graph is



(b) Is 0 in the range of $g(x)$? Justify your answer.

No b/c there is no value of x for which $b^x = 0$.

2. (4 points) Let $T(t)$ be the temperature of an object (in degrees Celsius) t hours after noon. What does the statement $T'(2.5) \approx -7$ mean in this context? Include units in your answer.

at 2.5 hours after noon (aka 2:30) the temperature is decreasing at a rate of 7°C/hr .

3. Suppose $g(x) = 0.3f(x) - 2$.

(a) (5 points) Describe how the graphs of $g(x)$ and $f(x)$ are related. (Use words like horizontal, vertical, compressed, shifted, stretched, translated, etc.)


first $f(x)$ is compressed vertically then shifted down 2 units to get $g(x)$.

(b) (4 points) Let $f'(x) = \sqrt{x^2 + 3}$. Evaluate $g'(1)$.

$$g'(x) = 0.3f'(x) - 0 = 0.3f'(x)$$

$$g'(1) = 0.3\sqrt{1+3} = 0.6$$

4. (5 points each) The graph below is a graph of $y = g''(x)$.



(a) Estimate $g'''(0)$. = slope of line tangent to $g''(x)$ at $x=0$

$$g'''(0) \approx \frac{2.5 - 4}{0 - (-1)} = \frac{-1.5}{1} = -1.5$$

(b) If possible, determine each of the following. Justify your answer.

i. The interval(s) for which g is decreasing.

To determine when g is decreasing we need to know when g' is negative but this can't be determined from the g'' graph.

ii. The interval(s) for which g is concave up.

g is concave up when $g''(x) > 0$
 so $(-3, 2) \cup (2, 3.8)$

Other acceptable answers


$$(-3, 2) \cup (2, 4)$$

$$(-3, 2) \cup (2, \infty)$$

$$(-3, 4)$$

$$(-3, \infty)$$

5. (4 points each) The graph of $f(x)$ is given. Solve the following (assume the tickmarks occur at 1, 2, etc).



(a) $\lim_{x \rightarrow -2^-} f(x)$ 1

(b) $\lim_{x \rightarrow -2^+} f(x)$ -2.5

(c) $f(-2)$ -2.5

(d) $\lim_{x \rightarrow 1} f(x)$ DNE b/c left and right hand limits do not agree

(e) $\lim_{x \rightarrow 3} f(x)$ -2

- (f) For what value(s) of x is $f(x)$ NOT continuous?

$$x = -3, -2, 1, 3$$

6. (6 points) Let $f(x) = \sqrt{x+2}$. Use the limit definition of derivative to compute $f'(14)$.

$$\begin{aligned} f'(14) &= \lim_{h \rightarrow 0} \frac{f(14+h) - f(14)}{h} = \lim_{h \rightarrow 0} \frac{\sqrt{14+h+2} - \sqrt{16}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{16+h} - 4}{h} \cdot \frac{(\sqrt{16+h} + 4)}{(\sqrt{16+h} + 4)} = \lim_{h \rightarrow 0} \frac{16+h - 16}{h(\sqrt{16+h} + 4)} \\ &= \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{16+h} + 4)} = \lim_{h \rightarrow 0} \frac{1}{\sqrt{16+h} + 4} = \frac{1}{\sqrt{16} + 4} = \boxed{\frac{1}{8}} \end{aligned}$$

7. (6 points each) Let $f(x) = 3x^2 + \frac{\pi}{x^2} - \sqrt{x} + 12$. = $3x^2 + \pi x^{-2} - x^{1/2} + 12$

(a) Find the derivative of f .

$$\begin{aligned} f' &= 3(2)x^1 + \pi(-2)x^{-3} - \left(\frac{1}{2}\right)x^{-1/2} + 0 \\ &= 6x - \frac{2\pi}{x^3} - \frac{1}{2}x^{-1/2} \end{aligned}$$


(b) Find an antiderivative of f .

$$\begin{aligned} F &= \frac{3x^3}{3} + \frac{\pi x^{-1}}{-1} - \frac{x^{3/2}}{3/2} + 12x + C \\ &= x^3 - \frac{\pi}{x} - \frac{2}{3}x^{3/2} + 12x + C \end{aligned}$$

8. (5 points each) Let $f(x)$ be a continuous function with the following properties:

- f is positive on the interval $(-\infty, 2)$ and negative on the interval $(2, \infty)$.
- $f'(x) < 0$ on the interval $(-\infty, 3)$, $f'(3) = 0$, and $f'(x) > 0$ on the interval $(3, \infty)$.
- $f''(x) > 0$ on the interval $(-\infty, 4)$, $f''(4) = 0$ and $f''(x) < 0$ on the interval $(4, \infty)$.

(a) Sketch a possible graph of $f(x)$.



(b) Suppose F is an antiderivative of f . At which value(s) of x in the interval $[0, 5]$ does F have a local minimum? Justify your answer.


None b/c F has a local minimum at a point where $F' = f$ changes from negative valued to positive valued, but that never happens.

(c) Suppose F is an antiderivative of f . Does F have an inflection point? Justify your answer.

Yes at $x = 3$ b/c $F'' = f'$ and f' changes sign at $x = 3$.

(d) Which value is larger: $\frac{f(0) - f(-3)}{3}$ or $f'(0)$? Justify your answer.

$\frac{f(0) - f(-3)}{3}$ is the slope of the secant line between the points $(0, f(0))$ and $(-3, f(-3))$.

On this region $f(x)$ is decreasing and concave up, like .

slope of dotted line = $\frac{f(0) - f(-3)}{3}$


slope of solid line = $f'(0)$

Both are negative but $f'(0)$ is larger (closer to 0).

9. (2 points) Who do you think will win the World Series?

- Atlanta Braves
- Baltimore Orioles
- Cincinnati Reds
- Detroit Tigers
- New York Yankees
- Oakland Athletics
- San Francisco Giants
- St. Louis Cardinals
- Texas Rangers
- Washington Nationals

Copyright 1996 Randy Glasbergen. www.glasbergen.com



**Many students actually look forward
to Mr. Atwadder's math tests.**