

Math 105: Review for Exam II - Solutions

1. Find dy/dx for each of the following.

(a) $y = x^2 + 2^x + e^2 + e^{2x} + \ln 2 + \ln(2x) + (\ln 2)x + \arctan 2$

$$\frac{dy}{dx} = 2x + (\ln 2)2^x + 2e^{2x} + \frac{1}{2x} \cdot 2 + \ln 2 \quad \text{Note that } e^2, \ln 2, \text{ and } \arctan 2 \text{ are constants.}$$

(b) $y = \sqrt{x} \cdot \arctan(5x)$

$$\frac{dy}{dx} = \frac{1}{2}x^{-1/2} \arctan(5x) + \sqrt{x} \cdot \frac{1}{1+(5x)^2} \cdot 5 = \frac{\arctan(5x)}{2x^{1/2}} + \frac{5\sqrt{x}}{1+25x^2}$$

(c) $y = \ln(\tan(2^{\cos(x^2)}))$

$$\frac{dy}{dx} = \frac{1}{\tan(2^{\cos(x^2)})} \cdot \sec^2(2^{\cos(x^2)}) \cdot \ln 2(2^{\cos(x^2)}) \cdot (-\sin(x^2)) \cdot 2x$$

(d) $y = \frac{x + e^\pi}{\cos 4 + \sin^5(6x)}$

Note that e^π and $\cos 4$ are constants.

$$\frac{dy}{dx} = \frac{(1)(\cos 4 + \sin^5(6x)) - (x + e^\pi)(5 \sin^4(6x) \cdot \cos(6x) \cdot 6)}{(\cos 4 + \sin^5(6x))^2} \quad \text{Recall that } \sin^5(6x) = (\sin(6x))^5.$$

(e) $y = (x^2 + 1)^{\sin x}$

We need logarithmic differentiation here.

$$\ln y = \sin x \cdot \ln(x^2 + 1)$$

Take natural log of each side.

$$\frac{1}{y} \frac{dy}{dx} = \cos x \cdot \ln(x^2 + 1) + \sin x \cdot \frac{1}{x^2 + 1} \cdot 2x$$

Differentiate.

$$\frac{dy}{dx} = \left[\cos x \cdot \ln(x^2 + 1) + \frac{2x \sin x}{x^2 + 1} \right] y$$

Solve for $\frac{dy}{dx}$.

$$\frac{dy}{dx} = \left[\cos x \cdot \ln(x^2 + 1) + \frac{2x \sin x}{x^2 + 1} \right] \cdot (x^2 + 1)^{\sin x}$$

Replace y .

2. Consider the curve defined by $x^3 + y^3 = \frac{9}{2}xy$ (known as the Folium of Descartes).

(a) Find dy/dx .

Use implicit differentiation.

$$\begin{aligned} 3x^2 + 3y^2 \frac{dy}{dx} &= \frac{9}{2}y + \frac{9}{2}x \frac{dy}{dx} \\ 3y^2 \frac{dy}{dx} - \frac{9}{2}x \frac{dy}{dx} &= \frac{9}{2}y - 3x^2 \\ \frac{dy}{dx} \left(3y^2 - \frac{9}{2}x \right) &= \frac{9}{2}y - 3x^2 \\ \frac{dy}{dx} &= \frac{\frac{9}{2}y - 3x^2}{3y^2 - \frac{9}{2}x} \end{aligned}$$

(b) Verify that the point (1,2) is on the curve above.

We must check to see if the values $x = 1$ and $y = 2$ satisfy the equation above.

$$\begin{aligned} x^3 + y^3 &\stackrel{?}{=} \frac{9}{2}xy \\ 1^3 + 2^3 &\stackrel{?}{=} \frac{9}{2} \cdot 1 \cdot 2 \\ 9 &\stackrel{?}{=} 9 \end{aligned}$$

Thus, the point (1,2) is on the curve.

- (c) **Find the equation of the tangent line at the point (1,2).**

We want $y = mx + b$.

$$m = \frac{\frac{9}{2} \cdot 2 - 3 \cdot 1^2}{3 \cdot 2^2 - \frac{9}{2} \cdot 1} = \frac{4}{5}, \text{ so } y = \frac{4}{5}x + b.$$

Now plug in $x = 1$ and $y = 2$ to find b .

$$2 = \frac{4}{5} \cdot 1 + b \Rightarrow \frac{6}{5} = b$$

Therefore, we have $y = \frac{4}{5}x + \frac{6}{5}$.

3. **Evaluate the following limits.**

Throughout this solution, the symbol \star will stand for whatever notation your instructor prefers for using L'Hopital's Rule on the indeterminate form $0/0$; this may be $\stackrel{0/0}{=}$ or $\stackrel{L'H}{=}$ or $\stackrel{H}{=}$ or $\stackrel{0/0}{\sim}$ or "has the form $\frac{0}{0}$, and so, by L'Hopital's Rule, is equal to" or something else. The symbol \heartsuit will serve the same purpose for the indeterminate forms ∞/∞ and $-\infty/\infty$.

(a) $\lim_{x \rightarrow 1} \frac{x^3 - 1}{7 - 7x} \star \lim_{x \rightarrow 1} \frac{3x^2}{-7} = \frac{3}{-7} = -\frac{3}{7}$

(b) $\lim_{x \rightarrow 0} \frac{1 - \cos(2x)}{3^x} = \frac{0}{1} = 0$ Can't use (and don't need) L'Hopital's Rule!

(c) $\lim_{x \rightarrow 0} \frac{1 - \cos(4x)}{5x^2} \star \lim_{x \rightarrow 0} \frac{4 \sin(4x)}{10x} \star \lim_{x \rightarrow 0} \frac{16 \cos(4x)}{10} = \frac{16}{10} = \frac{8}{5}$

(d) $\lim_{x \rightarrow \infty} \frac{x^2}{2^x} \heartsuit \lim_{x \rightarrow \infty} \frac{2x}{\ln 2 \cdot 2^x} \heartsuit \lim_{x \rightarrow \infty} \frac{2}{\ln 2 \cdot \ln 2 \cdot 2^x} = 0$

4. **Consider the function $f(x) = x^4 e^x$ with domain all real numbers.**

- (a) **Find the x -value(s) of all roots (x -intercepts) of f .**

The equation $x^4 e^x = 0$ means $x^4 = 0$ (that is, $x = 0$) or $e^x = 0$ (no solution), so the only root is at $x = 0$.

- (b) **Find the x - and y -value(s) of all critical points and identify each as a local max, local min, or neither.**

$$\begin{aligned} f'(x) &= 4x^3 e^x + x^4 e^x \\ 0 &= x^3 e^x (4 + x) \\ \Rightarrow x &= 0, -4 \end{aligned}$$

Note that e^x is never 0.

	$x < -4$	$-4 < x < 0$	$4 < x$
f'	positive	negative	positive
f	\nearrow	\searrow	\nearrow

y -values: $f(-4) = 256e^{-4} \approx 4.689$, $f(0) = 0$

So, f has a local maximum at $(-4, 256e^{-4})$ and a local minimum at $(0, 0)$.

- (c) **Find the x - and y -value(s) of all global extrema and identify each as a global max or global min.**

There is a global minimum at $(0, 0)$. There is no global maximum because as $x \rightarrow \infty$, $f(x) \rightarrow \infty$. Note that as $x \rightarrow -\infty$, $f(x) \rightarrow 0$. You can verify this by using L'Hopital's Rule on x^4/e^{-x} .

(d) Find the x -value(s) of all inflection points.

$$f''(x) = 12x^2e^x + 4x^3e^x + 4x^3e^x + x^4e^x \quad \text{Use Product Rule on each product in } f'(x) \text{ above.}$$

$$0 = e^x(x^4 + 8x^3 + 12x^2)$$

$$0 = e^x x^2(x^2 + 8x + 12)$$


$$0 = e^x x^2(x + 2)(x + 6)$$

$$\Rightarrow x = 0, -2, -6$$

	$x < -6$	$-6 < x < -2$	$-2 < x < 0$	$0 < x$
f''	positive	negative	positive	positive
f	concave up	concave down	concave up	concave up

So, the x -values of the inflection points of f are $x = -2$ and $x = -6$ but NOT $x = 0$.

(e) Sketch f .




5. How would your answers to the previous question change if the domain of f were $[-10, 10]$?

There would be a global maximum at $(10, 10^4e^{10})$. (And the graph would be restricted to $-10 \leq x \leq 10$).

6. Find an antiderivative of $y = \frac{5}{\sqrt{1-9x^2}} + x^3 + \cos(2x) + e^3$.

The answer is $\frac{5 \arcsin 3x}{3} + \frac{x^4}{4} + \frac{\sin 2x}{2} + e^3x + C$, where C is any constant.

7. You are planning to build a box-shaped aquarium with no top and with two square ends. Your budget is \$288. If the glass for the sides costs \$12 per square foot and the opaque material for the bottom costs \$3 per square foot, what dimensions will maximize the volume? Be sure to show how you know you have found the maximum.



Goal: Maximize volume

Objective function: volume = $V = x \cdot x \cdot y = x^2y$

We need to get this down to a function of just one variable, so we use the *constraint equation*:

$$\begin{aligned}\text{total cost} &= (\text{cost of base}) + (\text{cost of two square ends}) + (\text{cost of two other sides}) \\ 288 &= 3xy + 12 \cdot 2x^2 + 12 \cdot 2xy \\ 288 &= 27xy + 24x^2 \\ 288 - 24x^2 &= 27xy \\ \frac{288 - 24x^2}{27x} &= y\end{aligned}$$

Substituting this back into the objective function gives

$$V = x^2y = x^2 \cdot \frac{288 - 24x^2}{27x} = x \cdot \frac{288 - 24x^2}{27} = \frac{1}{27}(288x - 24x^3).$$

Now that we have V as a function of just one variable, we find its maximum.

$$\begin{aligned}V'(x) &= \frac{1}{27}(288 - 72x^2) \\ 0 &= \frac{1}{27}(288 - 72x^2) \\ 0 &= (288 - 72x^2) \\ 72x^2 &= 288 \\ x^2 &= \frac{288}{72} \\ x &= 2\end{aligned}$$

We discard $x = -2$ because lengths must be nonnegative.

Since V' is positive for $x < 2$ and negative for $2 < x$, we know that the maximum occurs at $x = 2$.

And $y = \frac{288 - 24x^2}{27x} = \frac{288 - 24 \cdot 2^2}{27 \cdot 2} = \frac{32}{9}$, so the dimensions are 2 by 2 by $\frac{32}{9}$.